



**THE PERMANENTE
MEDICAL GROUP**

SRO-MAIN CAMPUS
401 BICENTENNIAL WAY
SANTA ROSA CA 95403-
2149
Encounter Record

Hardeman, Edwin
MRN: [REDACTED], DOB: [REDACTED], Sex: M
Encounter date: 3/9/2006

Patient Email
3/9/2006

Visit Information

Date & Time	Provider	Department	Encounter #
3/9/2006	Ruffner-Statzer, Susan Marie (M.D.)	GASTROENTEROLOGY	870934096

Encounter Messages

(no subject)

From	To	Sent and Delivered
Susan Marie (M.D.) Ruffner-Statzer Last Read in kp.org 3/9/2006 3:46 PM by Edwin Hardeman	Edwin Hardeman	3/9/2006 1:45 PM

Ed, I got your message about WHAT DOES THE NEGATIVE VIRAL LOAD MEAN? What it means is that there is no detectable virus in your blood. It does not mean that the virus is completely gone from your liver. It does mean that the treatment appears to be working and that it is worthwhile to continue the treatment. We will test for the virus again at 24 weeks, at the end of the treatment and 6 months after completing the treatment. It is very important to continue to take your ribavirin faithfull twice a day, every day. Hope that helps. Dr. Ruffner

Patient Secure Message

(no subject)

From	To	Sent and Delivered
Susan Marie (M.D.) Ruffner-Statzer Last Read in kp.org 3/9/2006 3:46 PM by Edwin Hardeman	Edwin Hardeman	3/9/2006 1:45 PM

Ed, I got your message about WHAT DOES THE NEGATIVE VIRAL LOAD MEAN? What it means is that there is no detectable virus in your blood. It does not mean that the virus is completely gone from your liver. It does mean that the treatment appears to be working and that it is worthwhile to continue the treatment. We will test for the virus again at 24 weeks, at the end of the treatment and 6 months after completing the treatment. It is very important to continue to take your ribavirin faithfull twice a day, every day. Hope that helps. Dr. Ruffner

Vitals

None

Glaucoma	IOP	C/D ratio	CCT
OD			514
OS			515



Patient Preferred Languages

Interpreter Needed	Spoken Language	Written Language
No	English	English

All Meds and Administrations

(There are no med orders for this encounter)

Allergies as of 3/9/2006

Reviewed On: 2/13/2006 By: Powell, Rosanne M (M.A.)

No Known Drug Allergies	Noted	Reaction Type	Reactions
2001-11-27; ADMIT ORDER	11/27/2001		

Future Appointments 9/21/2018 - 9/21/2019

Generated on 9/21/18 11:10 AM
CONFIDENTIAL

HARDEMAN_EDWIN_KAISERSANTAROSA_00230

UNITED STATES DISTRICT COURT
NORTHERN DISTRICT OF CALIFORNIA
TRIAL EXHIBIT 67
Case No. 3:16-cv-0525-VC
Date Entered _____
By _____
Deputy Clerk



SRO-MAIN CAMPUS
 401 BICENTENNIAL WAY
 SANTA ROSA CA 95403-2149
 Encounter Record

Hardeman, Edwin
 MRN: [REDACTED], DOB: [REDACTED], Sex: M
 Encounter date: 3/9/2006

Patient Email
 3/9/2006

Visit Information

Date & Time	Provider	Department	Encounter #
3/9/2006	Ruffner-Statzer, Susan Marie (M.D.)	FAMILY MEDICINE SUITE 110	870934097

Encounter Messages

RE: Test

From	To	Sent
Edwin Hardeman	Susan Marie (M.D.) Ruffner-Statzer	3/10/2006 12:25 PM

Thank You, that is all I wanted to know.

Previous Messages

----- Message -----

From: SUSAN M. RUFFNER-STATZER MD
 Sent: 3/10/06 12:20 PM
 To: Edwin Hardeman
 Subject: RE: Test

Ed, There is no practical test for virus in the liver. The virus will IMMEDIATELY return if you stop taking the medication now. If you finish the 48 weeks of therapy and there is no detectable virus in your blood, you have a 50:50 chance that it will return in the next 6 months. There is not a lot of data about Long-term use of ribavirin (i.e. more than 48 weeks. If the virus stays undetectable after 6 months off treatment, you are likely CURED. We will continue to test your blood for return of virus, but 95% of the time (95/100) it stays gone. If it is gone after 5 years, we will call you cured. I hope that helps. The bottom line is "it's working" and "don't quit now:" Dr. Ruffner

----- Message -----

From: HARDEMAN, EDWIN
 Sent: 3/9/2006 4:51 PM
 To: Office of SUSAN M. RUFFNER-STATZER MD
 Subject: Test

Is there a test that can detect the any virus in my liver ? If there is no virus in my blood is taking the ribavirin safe to continue using long term ?

RE: Test

From	To	Sent and Delivered
Susan Marie (M.D.) Ruffner-Statzer Last Read in kp.org 3/10/2006 12:25 PM by Edwin Hardeman	Edwin Hardeman	3/10/2006 12:20 PM

Ed, There is no practical test for virus in the liver. The virus will IMMEDIATELY return if you stop taking the medication now. If you finish the 48 weeks of therapy and there is no detectable virus in your blood, you have a 50:50 chance that it will return in the next 6 months. There is not a lot of data about Long-term use of ribavirin (i.e. more than 48 weeks. If the virus stays undetectable after 6 months off treatment, you are likely CURED. We will continue to test your blood for return of virus, but 95% of the time (95/100) it stays gone. If it is gone after 5 years, we will call you cured. I hope that helps. The bottom line is "it's working" and "don't quit now:" Dr. Ruffner